TAXONOMY OF EURYTAENIA (APIACEAE)

Guy L. Nesom

2925 Hartwood Drive Fort Worth, Texas 76109 guynesom@sbcglobal.net

ABSTRACT

Eurytaenia comprises two species, E. texana and E. hinckleyi, essentially distinguished in morphology only by differences in fruit. Eurytaenia hinckleyi is known only from six counties in southwestern Texas and adjacent New Mexico, while E. texana is more widespread in Texas and western Oklahoma. A key, descriptions, distribution map, and illustrations of fruits and plants are provided.

KEY WORDS: Eurytaenia texana, Eurytaenia hinckleyi, Apiaceae, fruit morphology

Eurytaenia has interesting features of morphology and geography. Observations of a taxonomic study of Eurytaenia are put on record here in a synopsis of the genus.

EURYTAENIA Torrey & A. Gray, Fl. N. Amer. 1: 633. 1840. Type: Eurytaenia texana Torrey & A. Gray

Annual herbs, aromatic with parsley odor; slender-taprooted. Stems 3–12 dm, simple or branching mostly from the base, scabrous-papillate in the inflorescence or completely glabrous or glabrate. Leaves: basal 1-pinnate, segments lanceolate to ovate-lanceolate with coarsely serrate margins, cauline 2-pinnate (-3-pinnate) blades ovate in outline, ultimate segments linear to filiform with entire margins; glabrous. Peduncles terminal and lateral. Umbels once-compound; involucral bracts 3-cleft; involucel bracts entire or 3-cleft, distinctly scarious-margined toward the base. Flowers: peripheral and central similar, protandrous; sepals distinct, linear to triangular; petals white, margins entire, apex inflexed; stylopodium depressed-conic, nearly flat. Schizocarps splitting, ellipsoid or oblong-ellipsoid to broadly ellipsoid or suborbicular, strongly flattened laterally, dorsal ribs 3, filiform, lateral wings thickened and abruptly or gradually thinning, oil tubes large and flattened, solitary in the intervals, 2 on the commissural face, mericarp dorsal surfaces usually scabrous-papillate; carpophore divided completely to the base along the whole length. x = 7. Derivation of name: Greek, *eury*, wide, *tainia*, ribbon or band, alluding to the broad oil tubes, especially those on the commissural faces of E. *texana* mericarps.

In addition to the annual duration and distinctive fruit morphology (Figs. 1, 2), plants of *Eurytaenia* can easily be recognized by their leaf morphology (Figs. 4, 5). The basal are once-compound with the segments lanceolate to ovate-lanceolate and coarsely serrate margined. The cauline become 2-(-3)-pinnate and the segments much narrower with entire margins, the medial and distal with filiform to linear segments. All leaves are usually persistent and the transition in morphology usually is evident on a single plant.

As surely did Coulter & Rose, Mathias, and Constance, I searched for any other feature outside of mature fruit morphology that would distinguish the two species but found none. *Eurytaenia hinckley* is distinct in its mature mericarps that are relatively narrower in shape with relatively narrower bodies, thickened lateral wings, and a thin dermal covering over the dorsal oil tubes; the commissural oil tubes are narrower, differently shaped, and sparsely papillate (Fig. 1). The difference between the two species is remarkably parallel to the difference between *Polytaenia texana* and *P. nuttallii* (Nesom 2012).

Although the mericarp differences show only relatively late in ontogeny, it often is possible to see the developing distinction soon after anthesis. In Eurytaenia texana development of the lateral wings is simultaneous at all points from the base to apex. In Eurytaenia hinckleyi the thickening begins at the apex (at the mericarp shoulders) and proceeds basipetally.



Figure 1. Mericarps of Eurytaenia hinckleyi (A) and E. texana (B), dorsal and commissural faces.



Figure 2. Umbellule of schizocarps of Eurytaenia hinckleyi. From MO isotype.




Figure 3. Geographic distribution of Eurytaenia hinckleyi and E. texana. Symbols without inner dots are from literature records, vouchers not seen. See text (E. texana) for documentation of the counties in the two isolated clusters.

1. Mericarps broadly elliptic to suborbicular, thickened near the body but becoming thinner than the body toward the margins; oil tubes not covered by pericarp or epidermis, those of commissural face 1. Mericarps ellipsoid to oblong-ellipsoid, thickened to the abruptly narrowed-rounded margins, thicker than the body; oil tubes of both surfaces lightly covered by pericarp or epidermis and papillate, those of commissural face narrowly lenticular in outline Eurytaenia hinckleyi



Figure 4. Representative plant of Eurytaenia texama. Garza County, Texas.



Figure 5. Representative plant of Eurytaenia hinckleyi. Ward County, Texas.

1. Eurytaenia texana Torrey & A. Gray, Fl. N. Amer. 1: 633. 1840. TYPE: USA. Texas. Austin Co.: San Felipe, 1835, T. Drummond s.n. (holotype: GH).

Eurytaenia macrophylla Buckley, Proc. Acad. Nat. Sci. Philadelphia 13: 455. 1861. TYPE: USA. **Texas.** Washington Co.: no other collection data, ["Dr. Linsecom" fide protologue] Lincecum s.n. (holotype: PH).

Leaves: blades 4–10 cm x 2–5 cm, lobed or pinnatifid with obtuse, crenate, to serrate lobes, petioles 2-5 cm, cauline pinnately or 3-pinnately dissected, divisions oblong-lanceolate to linear or filiform, margins sharply serrate to entire, terminal often elongate. Peduncles 4–15 cm. Umbels: involucre of ca. 5 usually 3-cleft bracts 5–10 mm; umbellules 10–26, rays 20–80 mm, strongly unequal; flowers 10-22 per umbellule. Pedicels 4-8 mm. Mericarps broadly elliptic to suborbicular, 4–6 mm x 4–5 mm, scabrous-papillate dorsally, glabrous on the commissure, lateral wings often purplish, thickened near the body but becoming thinner than the body toward the margins; oil tubes not covered by pericarp or epidermis, dorsal usually papillate, commissural completely glabrous, those of commissural face depressed semi-circular in outline. 2n = 14 (Bell & Constance 1957).

Flowering (Mar-)Apr-Jul(-Aug). Sand and sandy loam, loose alluvial sand over granite, dune-like deposits, abandoned fields, pastures, roadsides, sandy prairies, mesquite savannas, floodplains, oak-juniper woodlands, live oak-post oak woodlands; (20–)50–900 m; Oklahoma, Texas.

There appears to be a hiatus in distribution of *Eurytaenia texana* between populations of the Coastal Bend area and those of central Texas and western Oklahoma (Fig. 2), and the habitats are generally different between the two areas. No differences in morphology are apparent, however, and all plants are identified as E. texana.

Documentation for the seeming disjunct cluster of three eastern counties is by the types of Eurytaenia texana (Austin Co.), E. macrophylla (Washington Co.), and a citation by Coulter and Rose (1900) — Hall 256 from Waller Co. (the specimen presumably at US). Records for Hood and Tarrant counties also are documented by citations from Coulter and Rose (1900) — Reverchon in June 1882 and Reverchon in June 1879, respectively.

2. Eurytaenia hinckleyi Mathias & Constance, Contr. Texas Res. Found., Bot. Stud. 1: 2. 1950. TYPE: USA. Texas. Andrews Co.: Shafter Lake, 10 Jul 1941, B.C. Tharp s.n. (holotype: MO digital image!; isotype: SMU!).

Leaves: blades 4–10 cm x 2–5 cm, lobed or pinnatifid with obtuse, crenate, to serrate lobes, petioles 2–5 cm, cauline pinnately or 3-pinnately dissected, divisions oblong-lanceolate to linear or filiform, margins sharply serrate to entire, terminal often elongate. Peduncles 4–15 cm. Umbels: involucral bracts ca. 5, 5–10 mm; umbellules 4–14, rays 12–45 mm, strongly unequal; flowers (4–)8–22 per umbellule. Pedicels 3–4 mm. Mericarps ellipsoid to oblong-ellipsoid, 5–8 mm x 4–5 mm, scabrous-papillate dorsally, mostly glabrous on the commissure, lateral wings not purplish, thickened to the abruptly narrowed-rounded margins, thicker than the body; oil tubes of both surfaces lightly covered by pericarp or epidermis and papillate, those of commissural face narrowly lenticular in outline. 2n = 14 (Bell & Constance 1957).

Flowering May-Jun(-Jul). Loose sand, sandy soil, dunes, openings in sandy mesquite woodland, sandy roadsides, less commonly gravelly soil of limestone hills; 800-1000 m; New Mexico (Chaves, Eddy, and Lea cos.), Texas (Andrews, Reeves, Ward, and Winkler cos.)

All collections of Eurytaenia hinckleyi except one have been made from habitats of loose sand — the record from Reeves Co. seems anomalous in habitat but it perhaps demonstrates an ecological breadth analogous to that of E. texana, as noted above. Reeves Co.: near Pecos, abundant in gravelly soil of limestone hills, 1 Jun 1932, Whitehouse 8371 (SMU).

LITERATURE CITED

- Bell, C.R. and L. Constance. 1957. Chromosome numbers in Umbelliferae. Amer. J. Bot 44: 565-572.
- Correll, D.S. and M.C. Johnston. 1970. Manual of the Vascular Plants of Texas. Texas Research Foundation, Renner, Texas. [Umbelliferae, pp. 1139–1169, by Mathias & Constance]
- Coulter and Rose. 1900. Monograph of the North American Umbelliferae. Contr. US Natl. Herb. 7: 9–422. Eurytaenia, pp. 126–127.
- Diggs, G.M., B.L. Lipscomb, and R.J. O'Kennon. 1999. Illustrated Flora of North Central Texas. Sida Bot. Misc. 16.
- Jones, S.D., J.K. Wipff, and P.M. Montgomery. 1997. Vascular Plants of Texas: A Comprehensive Checklist Including Synonymy, Bibliography, and Index. Univ. of Texas Press, Austin.
- Mathias, M.E. and L. Constance. 1945. Eurytaenia. Genus 80. N. Amer. Fl. 28B: 203–204.
- Mathias, M.E. and L. Constance. 1961 [issued separately in 1951]. Umbelliferae. Pp. 263–329, plates 1–54, in C.L. Lundell and collaborators. Flora of Texas, Vol. 3. Texas Research Foundation, Renner.
- Nesom, G.L. 2012. Taxonomy of *Polytaenia* (Apiaceae): P. nuttallii and P. texana. Phytoneuron 2012-n: 1–11.
- Oklahoma Vascular Plants Database (OVPD). 2012. Oklahoma Biological Survey, Biodiversity information and data. http://www.biosurvey.ou.edu/atlasdesc.html
- Turner, B.L., H. Nichols, G. Denny, and O. Doron. 2003. Atlas of the Vascular Plants of Texas. Vol. I–Dicots. Sida, Bot. Misc. 24.